1) The acid dissociation constant ( $K_a$ ) for benzoic acid is 6.3 x  $10^{-5}$ . Find the pH of a 0.35 M solution of benzoic acid.

2) Find the pH of a 0.275 M hypochlorous acid solution.  $K_a = 3.0 \times 10^{-8}$ .

3) Find the pH of a solution that contains 0.0925 M nitrous acid ( $K_a = 4.5 \times 10^{-4}$ ) and 0.139 M acetic acid ( $K_a = 1.8 \times 10^{-5}$ ).

| 1) |             | HC <sub>7</sub> H <sub>3</sub> O | <sub>2(aq)</sub> | $_{(aq)} + C_7 H_3 O_2^{-}_{(aq)}$ | aq) |
|----|-------------|----------------------------------|------------------|------------------------------------|-----|
|    | initial     | 0.35 M                           | 0 M              | 0 M                                |     |
|    | change      | - x M                            | + x M            | +x M                               |     |
|    | equilibrium | (0.35 – x) M                     | хМ               | x M                                |     |

Note that:  $(0.35 - x) M \approx 0.35 M so$ 

$$K_a = [H^+][C_7H_3O_2^-] = (x)(x) = (x)(x) = x^2 = 6.3 \times 10^{-5}$$
  
 $[HC_7H_3O_2]$  (0.35 - x) (0.35)

$$x^2 = (6.3 \times 10^{-5}) (0.35) = 2.205 \times 10^{-5}$$

$$x = 4.7 \times 10^{-3} M$$
  $x = moles/L formed$ 

$$pH = - \log (4.7 \times 10^{-3}) = 2.33$$

Note that:  $(0.275 - x) M \approx 0.275 M so$ 

$$K_a = [H^+][ClO^-] = (x)(x) = (x)(x) = (x)(x) = x^2 = 3.0 \times 10^{-8}$$
  
[HClO] (0.275 - x) (0.275)

$$x^2 = (3.0 \times 10^{-8}) (0.275) = 8.25 \times 10^{-9}$$
  
  $x = 9.08 \times 10^{-5}$  M

$$x = 9.08 \times 10^{-5} M$$

$$pH = -\log(9.08 \times 10^{-5}) = 4.042$$

## 3) First the amount of H<sup>+</sup> from each acid must be calculated.

|             | HNO <sub>2(a</sub> | <sub>aq)</sub> ≒ H⁺ | $(aq)$ + $NO_2(aq)$ |
|-------------|--------------------|---------------------|---------------------|
| initial     | 0.0925 M           | 0 M                 | 0 M                 |
| change      | - x M              | + x M               | +x M                |
| equilibrium | (0.0925 – x) M     | x M                 | x M                 |

Note that:  $(0.0925 - x) M \approx 0.0925 M so$ 

$$K_a = [H^+][NO_2^-] = (x)(x) = (x)(x) = x^2 = 4.5 \times 10^{-4}$$
  
 $[HNO_2]$  = (0.0925 - x) = (0.0925)

$$x^2 = (4.5 \times 10^{-4}) (0.0925) = 4.1625 \times 10^{-5}$$
  
 $x = 6.45 \times 10^{-3}$  M  $x = \text{moles/L formed}$ 

|             | $HC_2H_3O_{2(aq)} = \Box$ | $H^+_{(aq)}$ + | $C_2H_3O_2^{-1}$ |
|-------------|---------------------------|----------------|------------------|
| initial     | 0.139 M                   | 0 M            | 0 M              |
| change      | - x M                     | + x M          | +x M             |
| equilibrium | (0.139 – x) M             | хM             | хM               |

Note that:  $(0.139 - x) M \approx 0.139 M so$ 

$$K_{a} = \underbrace{[H^{+}][C_{2}H_{3}O_{2}^{-}]}_{[H C_{2}H_{3}O_{2}]} = \underbrace{(x)(x)}_{(0.139 - x)} = \underbrace{(x)(x)}_{(0.139)} = \underbrace{x^{2}}_{(0.139)} = 1.8 \times 10^{-5}$$

$$x^2 = (1.8 \times 10^{-5}) (0.139) = 2.502 \times 10^{-6}$$
  
  $x = 1.58 \times 10^{-3} M$ 

Then add the results together and use that value to find the pH.

$$6.45 \times 10^{-3} M + 1.58 \times 10^{-3} M = 8.03 \times 10^{-3} M$$

$$pH = - log (8.03 \times 10^{-3}) = 2.095$$